



GLAST

Science Objectives

see <http://glast.gsfc.nasa.gov/>

also

LAT: <http://www-glast.stanford.edu> and

<http://www-glast.slac.stanford.edu/>

GBM: <http://gammarray.msfc.nasa.gov/gbm/>

Steven Ritz

steven.m.ritz@nasa.gov

GLAST Deputy Project Scientist



Outline



- ▶ ***Why γ 's? Why this energy range? Techniques***
- ▶ ***Brief Tour of Science Topics***
- ▶ ***Science Requirements***
- ▶ ***Science Mission Elements***
- ▶ ***Summary***



Why study γ 's?

Gamma rays carry a wealth of information:


- γ rays do not interact much at their source: they offer a direct view into Nature's largest accelerators.
- similarly, the Universe is mainly transparent to γ rays: can probe cosmological volumes. Any opacity is energy-dependent.
- conversely, γ rays readily interact in detectors, with a clear signature.
- γ rays are neutral: no complications due to magnetic fields. Point directly back to sources, etc.

Two GLAST instruments:

LAT: 20 MeV – >300 GeV

GBM: 10 keV – 25 MeV

Launch: 2007




Large Area Telescope
(LAT)

GLAST Burst Monitor
(GBM)



Cosmic γ -ray Measurement Techniques



For $E_\gamma < \sim 50$ GeV, must detect above atmosphere (balloons, satellites)

For $E_\gamma > \sim 50$ GeV, information from showers penetrates to the ground (Cerenkov, air showers)

Photon interaction mechanisms:




Fig. 2: Photon cross-section σ in lead as a function of photon energy. The intensity of photons can be expressed as $I = I_0 \exp(-\sigma x)$, where x is the path length in radiation lengths. (Review of Particle Properties, April 1980 edition).







Gamma-ray Experiment Techniques



Space-based:

- use pair-conversion technique



Ground-Based:


- Airshower Cerenkov Telescopes (ACTs)



image the Cerenkov light from showers induced in the atmosphere. Examples: Whipple, CANGAROO, HEGRA, STACEE, CELESTE, CELESTE, VERITAS, MAGIC, HESS



- Extensive Air Shower Arrays (EAS)




Directly detect particles from the showers induced in the atmosphere. Example: MILAGRO



Gamma-ray Observatories



Complementary capabilities




angular resolution
duty cycle
area
field of view
energy resolution

ground-based	space-based
<u>ACT</u>	<u>EAS</u>
good	fair
low	high
large	large
small	large _{+can reorient}
good	good, w/ smaller systematic uncertainties

The next-generation ground-based and space-based experiments are well matched.


Gamma-ray Observatories





An Important Energy Band for Cosmology

Photons with $E > 10$ GeV are attenuated by the diffuse field of UV-Optical-IR extragalactic background light (EBL)



No significant attenuation below ~ 10 GeV.

only $e^{-\tau}$ of the original source flux reaches us

EBL over cosmological distances is probed by gammas in the 10-100 GeV range. Important science for GLAST!

In contrast, the TeV-IR attenuation results in a flux that may be limited to more local (or much brighter) sources.




A dominant factor in EBL models is the time of galaxy formation -- attenuation measurements can help distinguish models.



EGRET



The high energy gamma ray detector on the Compton Gamma Ray Observatory (20 MeV - ~20 GeV)





The Success of EGRET: Probing New Territory

History:

SAS-2, COSB (1970's-1980's) exploration phase: established galactic diffuse flux

EGRET (1990's) established field:

- ★ increased number of ID'd sources by large factor;
- ★ broadband measurements covering energy range ~ 20 MeV - ~ 20 GeV;
- ★ discovered many still-unidentified sources;
- ★ discovered surprisingly large number of Active Galactic Nuclei (AGN);
- ★ discovered multi-GeV emissions from gamma-ray bursts (GRBs);
- ★ discovered GeV emissions from the sun

GLAST LAT will explore the unexplored energy range above EGRET's reach, filling in the present gap in the energy spectrum, and will cover the very broad energy range ~ 20 MeV - >300 GeV with superior acceptance and resolution. Historically, opening new energy regimes has led to the discovery of totally unexpected new phenomena.



GLAST Science

GLAST will have a very broad menu that includes:

- ▶ ***Systems with supermassive black holes***
- ▶ ***Gamma-ray bursts (GRBs)***
- ▶ ***Pulsars***
- ▶ ***Solar physics***
- ▶ ***Origin of Cosmic Rays***
- ▶ ***Probing the era of galaxy formation***
- ▶ ***Discovery! Particle Dark Matter? Other relics from the Big Bang?
Testing Lorentz invariance. New source classes.***

Huge increment in capabilities.

***GLAST draws the interest of both the the High Energy Particle Physics
and High Energy Astrophysics communities.***




GLAST LAT High Energy Capabilities



- Huge FOV (~20% of sky)
- Broadband (4 decades in energy, including unexplored region > 10 GeV)
- Unprecedented PSF for gamma rays (factor > 3 better than EGRET for E>1 GeV)
- Large effective area (factor > 4 better than EGRET)
- **Results in factor > 30-100 improvement in sensitivity**
- No expendables → long mission without degradation



Features of the gamma-ray sky



EGRET all-sky survey (galactic coordinates) $E > 100$ MeV

diffuse extra-galactic background
(flux $\sim 1.5 \times 10^{-5} \text{ cm}^{-2} \text{ s}^{-1} \text{ sr}^{-1}$)

galactic diffuse (flux $\sim O(100)$ times larger)

high latitude (extra-galactic) point sources
(typical flux from EGRET sources $O(10^{-7} - 10^{-6}) \text{ cm}^{-2} \text{ s}^{-1}$)

galactic sources (pulsars, un-ID'd)

An essential characteristic: VARIABILITY in time!

➔ Field of view, and the ability to repoint, important for study of transients.




Sources



Third EGRET Catalog

$E > 100 \text{ MeV}$

EGRET 3rd
Catalog: 271
sources



- ◆ Active Galactic Nuclei
- Unidentified EGRET Sources
- Pulsars
- LMC
- Solar FLare




Sources



5 σ Sources from Simulated One Year All-sky Survey

LAT 1st Catalog:
>9000 sources possible



Results of one-year all-sky survey.
(Total: 9900 sources)


● AGN
● 3EG Catalog

● Galactic Halo
● Galactic Plane



Unidentified Sources


172 of the 271 sources in the EGRET 3rd catalog are “unidentified”



- Rosat or Einstein X-ray Source
- 1.4 GHz VLA Radio Source

EGRET source position error circles are $\sim 0.5^\circ$, resulting in counterpart confusion.

GLAST will provide much more accurate positions, with ~ 30 arcsec - ~ 5 arcmin localizations, depending on brightness.




Cygnus region (15x15 deg)



Diffuse Extra-galactic Background Radiation


Is it really isotropic (e.g., produced at an early epoch in intergalactic space) or an integrated flux from a large number of yet unresolved sources? GLAST has much higher sensitivity to weak sources, with better angular resolution.



E > 1 GeV Map

GLAST will bring alive the HE gamma-ray sky.


Simulated LAT One Year All-sky map (E > 100 MeV)



The origin of the diffuse extragalactic gamma-ray flux is a mystery. Either sources are there for GLAST to resolve (and study!), OR there is a truly diffuse flux from the very early universe.



Virgo Region ($E > 1$ GeV)



EGRET One-Year All-Sky Survey ($E > 100$ MeV)





Active Galactic Nuclei (AGN)



Active galaxies produce vast amounts of energy from a very compact central volume.

Prevailing idea: powered by accretion onto super-massive black holes ($10^6 - 10^{10}$ solar masses). Different phenomenology primarily due to the orientation with respect to us.

HST Image of M87 (1994)




Models include energetic (multi-TeV), highly-collimated, relativistic particle jets. High energy γ -rays emitted within a few degrees of jet axis. Mechanisms are speculative; γ -rays offer a direct probe.



AGN Shine Brightly in GLAST Energy Range

Power output of AGN is remarkable. Multi-GeV component can be dominant!

Estimated luminosity of 3C 279:
~ 10^{45} erg/s
corresponds to 10^{11}
times total solar
luminosity
just in γ -rays!! Large
variability within days.



Multiwavelength Spectrum of 3C 279

Sum all the power over the whole electromagnetic spectrum from all the stars of a typical galaxy: an AGN emits this amount of power in JUST γ rays from a very small volume!




AGN: What GLAST will do



EGRET has detected ~ 70 AGN.

Extrapolating, GLAST should expect to see dramatically more – many thousands:

- Allows a statistically accurate calculation of AGN contribution to the high energy diffuse extra-galactic background.
- Constrain acceleration and emission models. How do AGN work?
- Large acceptance and field of view allow relatively fast monitoring for variability over time -- correlate with other detectors at other wavelengths.
- Probe energy roll-offs with distance (light-light attenuation): info on era of galaxy formation.
- Long mission life to see weak sources and transients.




Joining the unique capabilities of GLAST with other detectors will provide a powerful tool.



AGN: Future Prospects

- ▶ ***Multiwavelength studies will continue to be the key to understanding how the engines work***
- ▶ ***Models: same population of HE electrons produces both components***






GLAST Probes the Optical-UV EBL

• Important advances offered by GLAST:

- (1) thousands of blazars - instead of peculiarities of individual sources, look for systematic effects vs redshift.
- (2) key energy range for cosmological distances (TeV-IR attenuation more local due to opacity).

• Effect is model-dependent (this is good):






Caveats


- How many blazars have intrinsic roll-offs in this energy range (10-100 GeV)? (An important question by itself for GLAST!) Again, power of statistics is the key.
- What if there is conspiratorial evolution in the intrinsic roll-off vs redshift? There may also be independent constraints (e.g., direct observation of integrated EBL).
- Must measure the redshifts for a large sample of these blazars!



Transients Sensitivity During All-sky Mode



EGRET Fluxes



During the all-sky survey, GLAST will have sufficient sensitivity after O(1) day to detect (5σ) the weakest EGRET sources.


*zenith-pointed
 ^"rocking" all-sky scan: alternating orbits point above/below the orbit plane



Gamma Ray Bursts

Bursts are isotropic and non-repeating (as far as we can tell):

2704 BATSE Gamma-Ray Bursts



⇒ important to view as much of the sky as possible

confirmed to be at cosmological distances.

some association with SNe.




Gamma Ray Bursts

The questions persist : **What are they and how do they work??**

=>measure closer to the intrinsic high-energy scale.

Already good indications of interesting behavior:

EGRET has detected very high energy emission associated with bursts, including a 20 GeV photon ~75 minutes after the start of a burst:



+ Milagrito
evidence for
TeV emission
from GRB
970417 ApJ
533(2000)533.

Hurley et al., 1994


Future Prospects: GLAST will provide definitive information about the high energy behavior of bursts: LAT and GBM together will measure emission over >7 decades of energy.
Place your bets on additional TeV burst detections!



GRB941017




Recent analysis by Gonzalez et al., published in Nature



Compare data from EGRET and BATSE: Distinct high-energy component has different time behavior. What is the high-energy break and total luminosity?

Need GLAST data!


Learn important lessons from the past.





GRBs and Instrument Deadtime

Distribution for the 20th brightest burst in a year (Norris et al)



LAT will open a wide window on the study of the high energy behavior of bursts.


Time between consecutive arriving photons



Roles of the GBM

- ▶ ***provides spectra for bursts from 10 keV to 30 MeV, connecting frontier LAT high-energy measurements with more familiar energy domain;***

Simulated GBM and LAT response to time-integrated flux from bright GRB 940217
 Spectral model parameters from CGRO wide-band fit
 1 NaI (14 °) and 1 BGO (30 °)



- ▶ ***provides wide sky coverage (8 sr) -- enables autonomous repoint requests for exceptionally bright bursts that occur outside LAT FOV for high-energy afterglow studies (an important question from EGRET);***
- ▶ ***provides burst alerts to the ground.***



Burst Handling

- ▶ ***Upon detection of a gamma-ray burst (GRB), GLAST can take two DIFFERENT kinds of actions:***
 - *alerts to the ground*
 - *GBM and LAT can issue alerts independently*
 - *autonomous repoint to follow burst*
 - *both GBM and LAT can generate a repoint request*
 - *GBM will probably detect more bursts than LAT. However, if LAT detects a burst, it will probably provide a better localization of the burst position.*
 - *if both GBM and LAT generate a repoint request, it is a science issue (not a spacecraft issue) to choose the target repoint location. The GBM repoint request is therefore routed through LAT; LAT either passes the GBM request through or sends LAT's position.*
 - *there is never more than one repoint request per burst. The instrument FOVs are very large.*
 - *alerts happen much more frequently than repoint requests*



GRB Science Repoint Scenarios



Summary


During all-sky scanning operations, detection of a sufficiently bright burst will cause the observatory to interrupt the scanning operation autonomously and to remain pointed at the burst region during the non-occulted viewing time for a period of 5 hours (adjustable). There are two cases:

- 1. The burst occurs within the LAT FOV.** The observatory will slew to keep the burst direction within 30 degrees (adjustable) of the Z axis during the non-occulted viewing period (neglecting SAA effects). **Such events are likely to occur a few times per month.**
- 2. The burst occurs outside the LAT FOV.** If the burst is exceptionally bright, the observatory will slew to bring the burst direction within 30 degrees (adjustable) of the Z axis during the non-occulted viewing period (neglecting SAA effects). **Such events are likely to occur a few times per year.**





Particle Dark Matter

Some important models in particle physics could also solve the dark matter problem in astrophysics. If correct, these new particle interactions could produce an anomalous flux of gamma rays.



Just an example of what might be waiting for us to find!





Testing Lorentz Invariance

- ▶ **Lorentz Invariance breaking models can lead to different maximum velocities by particle type (Stecker&Glashow, Coleman&Glashow):**

$$c_e \equiv c_\gamma (1 + \delta), \quad 0 < |\delta| \ll 1$$

Stecker:

- For $\delta < 0$, photons can decay to e^+e^- pairs if $E_\gamma > m_e \sqrt{2/|\delta|}$
Observations of the Crab ($E > 50$ TeV) implies $-\delta < 2 \times 10^{-16}$
- For $\delta > 0$, superluminal electrons will emit vacuum Cerenkov radiation and the threshold for pair creation will be altered. Cosmic ray data and inferred information from Mrk501 blazar observations $\Rightarrow \delta < 3 \times 10^{-14} - 1.3 \times 10^{-15}$.

- ▶ **Some classes of QG models imply a linear photon velocity dispersion (Amelino-Camelia et al., Ellis, Mavromatos, Nanopoulos):**

$$V = c \left(1 - \xi \cdot \frac{E}{E_{QG}} + \dots \right)$$

Use GRBs! Effects could be $O(100)$ ms or larger, using GLAST data alone.

But ?? effects intrinsic to bursts?? Representative of window opened by measurements at such large distance and energy scales.



Requirements



- ▶ ***Mission Level II science requirements are codified in the Science Requirements Document (SRD)***
- ▶ ***See <http://glast.gsfc.nasa.gov/project/cm/mcdl/>***



Observatory Science Requirements (I)



	<i>Quantity</i>	<i>GLAST Requirement ¹</i>	<i>GLAST Goal ¹</i>	<i>GLAST Minimum ¹</i>	<i>Science Topic</i>
28	Mission Lifetime (<i><20% degradation</i>) ²	> 5 years	> 10 years	> 2 years	ALL
29	Telemetry Downlink Orbit Average	> 300 kbps	> 1 Mbps	> 300 kbps	ALL
30	Telemetry Downlink Realtime ³	> 1 kbps	> 2 kbps	> 0.5 kbps	GRBs
31	Telemetry Uplink Realtime ³	> 1 kbps	> 2 kbps	> 0.5 kbps	GRBs, AGN
32	Time to Respond to TOO's on Ground ⁴	< 6 hours	< 4 hours	< 12 hours	GRBs, AGN
33	Spacecraft Repointing Times for Autonomous Slews ⁵	< 10 min	< 5 min	NA	GRBs, AGN
34	GRB Notification Time to Ground by Spacecraft ⁶	< 7 sec	< 4 sec	< 10 sec	GRBs, AGN
35	Pointing Accuracy Absolute ⁷	< 2°	< 0.5°	< 5°	ALL



Observatory Science Requirements (II)



36	Pointing Knowledge ⁷	< 10 arcsec	< 5 arcsec	< 20 arcsec	ALL
37	Observing Modes	- Rocking zenith pointing ¹⁵ - Pointed mode ⁸			ALL
38	Targeting	No restrictions on Pointing ⁸ of axis normal to LAT			ALL
39	Uniformity of Sky Coverage during All-sky Survey ⁹	< ± 20%	< ± 10%	< ± 30%	ALL
40	Observatory Time Accuracy ¹⁰	< 10 μsec	< 3 μsec	< 30 μsec	Pulsars
41	Observatory Absolute Position Accuracy	< 3.3 km	< 1 km	< 10 km	Pulsars
42	Observing Efficiency ¹¹	> 90 %	> 95%	> 80%	ALL
43	Data Loss ¹²	< 2 %	< 1%	< 5%	ALL
44	Data Corruption ¹³	< 10 ⁻¹⁰	< 3 x 10 ⁻¹¹	< 3 x 10 ⁻¹⁰	ALL
45	Earth Avoidance ¹⁴	> 30°(adjustable)	N/A	N/A	ALL



Observatory Science Requirements (III)




- 1 Requirement = value to design to; Goal = value to strive for to enhance science; Minimum = value that if not satisfied triggers a Project review.
- 2 20% degradation = no more than 20% loss of LAT science return.
- 3 Uplink telemetry rate for at least 80% of time outside of SAA.
- 4 Response time for the SSC and MOC to plan and send a spacecraft repointing command after the decision is made to respond to a Target of Opportunity (TOO).
- 5 Time for 75° slew. note: being modified for single failed wheel case
- 6 Time from spacecraft receipt of GRB notification from GBM or LAT to delivery to the Gamma-ray Coordinates Network (GCN) computer for 80% of all GRBs detected by the GBM or LAT.
- 7 1 sigma radius.
- 8 Pointing of axis normal to LAT to within 30° of source. (No science constraint on roll axis.)
- 9 Sky coverage exposure uniformity integrating for 7 days, not including SAA effects.
- 10 Relative to Universal Time, 1 sigma r.m.s..
- 11 Fraction of time with data return, not including SAA effects.
- 12 Fraction of data taken by the instruments but not delivered to the IOC. Not including SAA data loss. Not including instrument downtime.
- 13 Fraction of undetected corrupted events.
- 14 Axis normal to LAT shall be capable of remaining at an angle greater than this above the Earth's horizon during normal operations, with the possible exception of rapid slewing to acquire a GRB.
- 15 The observatory shall be capable of scanning the LAT FOV (55° half-angle) over >90% of the celestial sphere repetitively on selectable timescales as short as every 2 orbits.



LAT Science Requirements Summary(I)

	Quantity	EGRET	LAT Requirement ¹	LAT Goal ¹	LAT Minimum ¹	Science Topic
1	Energy Range Low Limit	20 MeV	< 20 MeV	< 10 MeV	< 30 MeV	ALL
2	Energy Range High Limit	30 GeV	> 300 GeV	> 500 GeV	> 100 GeV	ALL
3	Effective Area ²	1500 cm ²	> 8000 cm ²	> 12,000 cm ²	> 8000 cm ²	ALL
4	Energy Resolution ³ (on-axis, 100 MeV - 10 GeV)	10%	< 10%	< 8%	< 20%	ALL
5	Energy Resolution ³ (on-axis, 10-300 GeV)		<20%	<15%	<30%	ALL
6	Energy Resolution (>60° incidence, >10 GeV) ⁴		< 6%	< 3%	NA ⁵	Dark Matter
7	Single Photon Angular Resolution - 68% ⁶ (on-axis, E>10 GeV)	0.5°	< 0.15°	< 0.1°	< 0.3°	ALL



LAT Science Requirements Summary(II)

8	Single Photon Angular Resolution - 68%⁶ (on-axis, E=100 MeV)	5.8°	< 3.5°	< 3°	< 5°	ALL
9	Single Photon Angular Resolution - 95%⁶ (on-axis)		< 3 x $\theta_{68\%}$	< 2 x $\theta_{68\%}$	< 4 x $\theta_{68\%}$	ALL
10	Single Photon Angular Resolution (off axis at 55°)		< 1.7 times on-axis	< 1.5 times on-axis	< 2 times on-axis	ALL
11	Field of View⁷	0.5 sr	> 2 sr	> 3 sr	> 1.5 sr	ALL
12	Source Location^{8,9} Determination	5 arcmin	< 0.5 arcmin	< 0.3 arcmin	< 1 arcmin	UGOs , GRBs
13	Point Source Sensitivity^{9,10} (> 100 MeV)	$\sim 1 \times 10^{-7}$ $\text{cm}^{-2} \text{s}^{-1}$	< $6 \times 10^{-9} \text{cm}^{-2} \text{s}^{-1}$	< $3 \times 10^{-9} \text{cm}^{-2} \text{s}^{-1}$	< $8 \times 10^{-9} \text{cm}^{-2} \text{s}^{-1}$	AGN, UGOs, Pulsars, GRBs
14	Instrument Time Accuracy¹¹	0.1 ms	< 10 μsec	< 2 μsec	< 30 μsec	Pulsars, GRBs
15	Background Rejection¹² (Contamination of high latitude diffuse sample in any decade of energy for >100 MeV.)	<1%	<10%	<1%	<15%	Diffuse
16	Dead Time	100 ms /event	< 100 μs /event	< 20 μs /event	< 200 μs /event	GRBs



LAT Science Requirements Summary(III)

17	GRB Location Accuracy On-Board ¹³		< 10 arcmin	< 3 arcmin	NA ⁵	GRBs
18	GRB Notification Time To Spacecraft ¹⁴		< 5 sec	< 2 sec	NA ⁵	GRBs

- 1 Requirement = value to design to; Goal = value to strive for to enhance science; Minimum = value that if not satisfied triggers a Project review.
- 2 Maximum (as function of energy) effective area at normal incidence. Includes inefficiencies necessary to achieve required background rejection. Effective area peak is typically in the 1 to 10 GeV range.
- 3 Equivalent Gaussian 1 sigma, on-axis.
- 4 Effective area for side incidence is 0.1 to 0.2 that of normal incidence for high resolution measurements.
- 5 NA = Not Applicable. Minimum values are not applicable for parameters that were not Requirements in the AO 99-OSS-03 Announcement of Opportunity.
- 6 Space angle.
- 7 Integral of effective area over solid angle divided by peak effective area. Geometric factor is Field of View times Effective Area.
- 8 High latitude source of $10^{-7} \text{ cm}^{-2} \text{ s}^{-1}$ flux at >100 MeV with a photon spectral index of -2.0 above a flat background and assuming no spectral cut-off. 1 sigma radius. 1-year survey.
- 9 Derived quantities delimited by double-lined box.
- 10 Sensitivity at high latitudes after a 1-year survey for a 5 sigma detection.
- 11 Relative to spacecraft time.
- 12 Assuming a high-latitude diffuse flux of $1.5 \times 10^{-5} \text{ cm}^{-2} \text{ s}^{-1} \text{ sr}^{-1}$ (>100 MeV) assuming a photon spectral index of -2.1 with no spectral cut-off.
- 13 For burst (< 20 sec duration) with > 100 photons above 1 GeV. This corresponds to a burst of $\sim 5 \text{ cm}^{-2} \text{ s}^{-1}$ peak rate in the 50 - 300 keV band assuming a spectrum of broken power law at 200 keV from photon index of -0.9 to -2.0. Such bursts are expected to occur in the LAT FOV ~ 10 times per year.
- 14 Time relative to detection of GRB.



GBM Science Requirements Summary (I)



	Quantity	BATSE	GBM Requirement ¹	GBM Goal ¹	GBM Minimum ¹	Science Topic
19	Energy Range Low Limit	25 keV	< 10 keV	< 5 keV	< 20 keV	ALL
20	Energy Range High Limit	10 MeV	> 25 MeV	> 30 MeV	> 20 MeV	ALL
21	Field of View ²	4 π	> 8 sr	> 10 sr	> 6 sr	ALL
22	Energy Resolution ³ (0.1 - 1.0 MeV)		< 10%	< 7%	< 12%	GRBs
23	GRB Alert Location ⁵		NA ⁴	< 15 deg	NA ⁴	GRBs
24	GRB Notification Time To Spacecraft ⁶		< 2 sec	< 1 sec	< 5 sec	GRBs
25	Dead Time Average		< 10 μ sec/event	< 3 μ sec/event	< 50 μ sec/event	GRBs
26	Instrument Time Accuracy ⁷	10 μ sec	< 10 μ sec	< 2 μ sec	< 30 μ sec	GRBs
27	Burst Sensitivity Ground Analysis ⁸	0.2 cm ⁻² s ⁻¹	< 0.5 cm ⁻² s ⁻¹	< 0.3 cm ⁻² s ⁻¹	< 1.0 cm ⁻² s ⁻¹	GRBs



GBM Science Requirements Summary (II)



46	Burst Sensitivity On-Board Trigger ⁹	0.3 cm ⁻² s ⁻¹	< 1.0 cm ⁻² s ⁻¹	< 0.75 cm ⁻² s ⁻¹	< 2.0 cm ⁻² s ⁻¹	GRBs
----	--	--------------------------------------	--	---	--	------

- 1 Requirement = value to design to; Goal = value to strive for to enhance science; Minimum = value that if not satisfied triggers a Project review.
- 2 Integral of effective area over solid angle divided by peak effective area. Geometric factor is Field of View times Effective Area. Should overlap with LAT FOV.
- 3 Equivalent Gaussian. 1 sigma. On axis.
- 4 NA= Not Applicable. The addition of the GRB monitor was a "goal" in the AO 99-OSS-03. The broad-band spectroscopic capability of the GRB instrument is upgraded here to be a requirement. The location of the bursts is listed only as a goal.
- 5 1 sigma radius. For burst of brightness 10 cm⁻² s⁻¹ in 50 - 300 keV band and a duration of 1 second or longer.
- 6 Time relative to a GBM GRB trigger. Used for both 'rapid ground notification' or 'burst alert' through TDRSS (or equivalent real-time link) and for 'LAT notification'.
- 7 Relative to spacecraft time.
- 8 GRB peak brightness sensitivity, 50 - 300 keV range, 5 sigma detection.
- 9 50% efficiency level for bursts occurring within the GBM FOV, excluding observational inefficiencies such as SAA passages and earth occultations, 50 - 300 keV range.



Science Mission Elements



▶ **Science Working Group (SWG)**

- *Membership includes the Interdisciplinary Scientists (IDS) and instrument team delegates.*
- *Bi-monthly telecons and ~bi-annual sit-down meetings, along with science symposia to engage the community. Minutes on the web.*

▶ **Users Committee**

- *independent of the SWG. External review/feedback on science tools planning and progress.*
- *includes members from both the astrophysics and high-energy particle physics communities who are likely users of GLAST data.*

▶ **GLAST Science Support Center (GSSC)**

- *Located at Goddard. Supports guest observer program, provides training workshops, provides data and software to community, archives through to HEASARC, joint software development with Instrument Teams.*



Operations Phases, Guest Observers, Data

- ▶ ***After the initial on-orbit checkout, verification, and calibrations, the first year of science operations will be an all-sky survey.***
 - *first year data used for detailed instrument characterization, refinement of the alignment, and key projects (source catalog, diffuse background models, etc.) needed by the community*
 - *data on transients will be released, with caveats*
 - *repoints for bright bursts and burst alerts are enabled*
 - *extraordinary ToO's supported*
 - *limited guest observer program*
 - *workshops for guest observers on science tools and mission characteristics for proposal preparation*

- ▶ ***Observing plan in subsequent years driven by guest observer proposal selections by peer review. All data released through the science support center (GSSC).***



Summary

GLAST will address many important questions:

- What is going on around black holes? How do Nature's most powerful accelerators work?*
- What are the unidentified sources found by EGRET?*
- What is the origin of the diffuse background?*
- What is the origin of cosmic rays?*
- What is the high energy behavior of gamma ray bursts?*
- When did galaxies form?*
- What else out there is shining gamma rays? Are there high-energy relics from the Big Bang? Are there further surprises in the poorly measured energy region?*

- ▶ *Large menu of “bread and butter” science, and large discovery potential.***
- ▶ *Science requirements are mature and stable.***
- ▶ *GLAST is part of the bigger picture of experiments at the interface between particle physics and astrophysics.***

We expect the gamma-ray community to grow in the GLAST era!!